



Agriculture and
Agri-Food Canada

Agriculture et
Agroalimentaire Canada

Unclassified / Non classifié

Canada

Weed management in potato production

Andrew McKenzie-Gopsill, PhD

Research Scientist – Weed Science

Agriculture and Agri-Food Canada, Charlottetown

andrew.mckenzie-gopsill@agr.gc.ca (902) 314-3683

© His Majesty the King in Right of Canada, as represented by the Minister of Agriculture and Agri-Food Canada (2024)

Weed competition costs Canadian potato producers upwards of \$83 million in lost yield potential annually



Weeds reduce potato yield by 44%




Minimal herbicide options



Reduced cultivation

Grasses vs Broadleaves

Weeds can be divided into two large groups




Monocots
Grasses



Dicots
Broadleaves

Identifying Grasses

Parts of a grass plant





© 2004 NC State University

Hairy

Ligules



CNAS Research & Extension, University of Gua

Absent



Membranous

Auricles



Vegetative Growth



Rhizomes
Under-ground stems




Tillers
Branches

Identifying Broadleaves



Broadleaf Morphology

Figure 1. Plant description.

1. Hypocotyl
2. Radicle
3. Cotyledon (simple leaf)
4. Cotyledonary node
5. Tap root
6. Lateral (branch) root
7. First true leaf (unifoliolate)
8. Trifoliolate leaflet
9. Terminal bud
10. Axillary buds
11. Hypocotyl arch
12. Nodes (point of leaf attachment)
13. Nodules
14. Root hairs



Cotyledon morphology



Leaf morphology



Compound

Leaf architecture



Opposite



Alternate



Rosette



Whorled

Vegetative reproduction



Stolons
Above-ground stems



Tubers- storage organs



Rhizomes
Below-ground stems




Adventitious root system

Other distinguishing features



Clasping leaves



Ocrea




Spines or thorns



Presence or absence of hair

Problem weeds of PEI



Annual Grasses




Barnyardgrass
Echinochloa crus-galli
No auricles
No ligule



Foxtails
Setaria spp.
Hairy ligule
Distinctive seed head



Annual Grasses



Crabgrasses (*Digitaria spp.*)

Membranous ligule

Smooth crabgrass may have hairs

Large crabgrass hairy on both leaf surfaces

Perennial Grasses



Quackgrass or Couchgrass
(*Agropyron repens*)
Membranous ligule
Clasping auricles
Rhizomes
Hair may be present or absent

Yellow nutsedge

Cyperus es

Triangula

Spreads by seed
and tuber



Joel Felix



Chickweeds (*Caryophyllaceae*)



Mouse-ear chickweed
Cerastium fontanum



Stitchwort
Stellaria graminea



Common chickweed
Stellaria media

Corn Spurry (*Caryophyllaceae*)



Mint Family (*Lamiaceae*)



Hemp-nettle (*Galeopsis tetrahit*)



Field mint (*Mentha arvensis*)



Bedstraw, Cleavers (*Galium spp.*)



Lambsquarters & Pigweed



Lambsquarters (*Chenopodium album*)

Redroot pigweed (*Amaranthus retroflexus*)

Mustard Family (*Brassicaceae*)



Wild mustard *Sinapis arvensis* Shepard's Purse *Capsella bursa-pastoris*

Stinkweed *Thlaspi arvense* Brown mustard *Brassica juncea*



Smartweed Family (*Polygonaceae*)



Wild buckwheat
Polygonum convolvulus



Lady's thumb
Persicaria maculosa




Smartweed
Polygonum spp.



Sheep sorrel
Rumex acetosella

Ragweed (*Ambrosia artemisiifolia*)



Common Mugwort (Asteraceae)



Sow Thistles (*Asteraceae*)



Spiny annual sow thistle
Sonchus asper

Perennial sow thistle
Sonchus arvensis



Annual sow thistle
Sonchus oleraceus

Goldenrod & Canada thistle



Canada goldenrod
Solidago canadensis



Canada thistle
Cirsium arvense

Less common



Iowa State University

Eastern black nightshade
Solanum ptychanthum



OMAFRA

Field violet
Viola arvensis



Wikipedia

Sand spurry
Spergularia rubra



MSU Turf

Corn speedwell
Veronica arvensis



Tufted vetch
Vicia cracca



© Peter Smith

Green pigweed
Amaranthus powellii




Wikipedia

Oxeye daisy
Leucanthemum vulgare

Volunteer Crops







Nearly half of potato fields surveyed have an herbicide resistant weed






Reduced
cultivation



Shorter crop
rotations



Reliance on
single modes of
action



Herbicide
Resistance

Scouting for Herbicide Resistance

Has the same herbicide mode of action been used repeatedly on this field?

Did the same rate previously control this weed?

Are there live and dead weeds following application?

Noticed a recent decline in control?

Are other weed species being controlled by this rate?

Spotting signs of PSII-inhibitor resistance



If resistance is suspected switch up your pre-emerge



Reflex



Frontier Max



Boundary LQD

Pre-emergent herbicides

In general can be applied at planting/ hilling all the way up to ground crack with no adverse affects on potato yield or quality

- Sencor
- Lorox
- Dual II Magnum
- Boundary LQD

- Glufosinate
- Glyphosate
- Frontier Max
- Reflex
- ~~Sencor STZ~~

Dual II Magnum (s-metolachlor)

- Group 15 – Inhibition of cell division
- Use rates between 1.25 and 1.75 L ha⁻¹
- Can apply from pre-planting all the way up to ground crack
- ***Has variety issues*** - potato variety responses not well studied
- Pre-emergent weed control
- Uptake in grasses through ***germinating shoot***, absorbed in ***shoots and roots of germinating broadleaves***
- Controls annual grasses and some broadleaf weeds (nightshade, suppression of pigweed)
- ***Rainfall within 10 days is required*** for maximum pre-emergence activity
- Can be soil incorporated to improve pre-emergent activity
- Provides 10-14 weeks of residual control
- Winter cereals can be planted 4-5 months post-application

Boundary LQD (s-metolachlor + metribuzin)

- Group 5 + Group 15
- Use rates of Boundary LQD between 1.85 and 2.5 L ha⁻¹
- Refer to label of Sencor and Dual if tank-mixing
- Broad spectrum of annual grass and broadleaf control
- ***Will not*** control PSII-inhibitor resistant weeds
- Resistance to s-metolachlor documented in pigweed species in the US

Results from our trials (Dual II Magnum)

Poor control of

- Lambsquarters

Moderate control of

- Redroot pigweed
- Shepard's purse
- Wild buckwheat & smartweeds
- Smooth crabgrass

Marketable yield equivalent to weed-free



Results from our trials (Boundary LQD)

Minimal control of PSII-inhibitor resistant species

Excellent control of all other species

See late season grasses

Marketable yield equivalent to weed-free



Potato cultivars vary in their tolerance to Dual II Magnum



Satina



Eva



Dakota



Red Norland



Payette



Targhee




Kennebec



Superior



Mountain Gem



Frontier Max (dimethenamid-P)

- Group 15 – Inhibition of cell division
- Use lower rates with reduced OM (OM~3% use 756 g ha⁻¹, consult label)
- Applied anytime after planting as long as **potatoes have not emerged**
- **Pre-emergent** control
- **Uptake via shoots and roots** but primarily through developing **coleoptile**
- Effective on annual grasses and some broadleaf weeds (pigweed, nightshade)
- Provides **season-long** weed control
- **Rainfall is required** within 7-10 days
- Can use shallow cultivation in absence of rainfall to move product to moist soil zone

Frontier Max (dimethenamid-P)

- In cold, wet conditions, potato emergence may be delayed or stunted
- If crop fails, ***do not*** re-plant potato – plant corn, soybean or dry bean
- Only ***a single application*** may be made per year
- Do not apply within 40 days of harvest
- Re-cropping intervals –
 - 100 days for cereals
 - Next spring for potato, corn, soybean, dry bean, cabbage
 - 11 months for all other crops
- 2016 resistance documented in pigweed species in Illinois
- Consult the label before using

Results from our trials

Poor control of

- Lambsquarters
- Shepard's purse
- Wild buckwheat & smartweeds
- Smooth crabgrass

Small marketable yield boost over weed-free (~10%)



Reflex (fomesafen)

- Group 14 – Inhibition of protoporphyrinogen oxidase (PPO)
- Use 1.0 L ha⁻¹, if weeds are emerged add Agral 90 @ 0.1% v/v
- Can be applied at planting up to potato emergence
- Provides *pre-* and *post-*emergent control of pigweed and suppression of lamb's quarters
- *Post-emergent* control of ragweed, wild mustard, lady's-thumb, Eastern black nightshade, canola
- For post-emergent control – max of 4 leaves for most weeds, 3 for lamb's quarters

Reflex (fomesafen)

- ***Do not*** cultivate within 7 days after application
- ***Do not*** apply to soils >5% OM or fine textured soils
- ***Do not*** apply Reflex to a field more than once every two years
- ***Do not*** harvest potatoes within 70 days of application
- Resistance documented in pigweed species across the soybean/corn belt of US
- Consult the label for using

Results from our trials

Good control of early emerging weeds
but doesn't get that second flush

We noted,

- Lambsquarters
- Smooth crabgrass
- Field mint
- Common chickweed

Marketable yield equivalent to weed-free



A single inter-row cultivation pre-hilling can reduce broadleaf weed biomass

Treatment	Weeks after planting	Lambsquarters (g m ⁻²)	Wild buckwheat (g m ⁻²)	Corn spurrey (g m ⁻²)	Yield (Cwt ac ⁻¹)
Single Cultivation	Three	4	19	9	312
Two Cultivations	Four	4	12	15	286
Three Cultivations	Five	8	28	19	277
Hilling only	Six	24	37	54	268
Boundary LQD	One	0	0.3	0	330

There are limited post-emergent chemical options available in potato



Fall & spring burndowns
can be used to control
perennials & winter
annuals prior to potato




Grass Herbicides

- Group 1 – ACCase inhibitor
- Post-emergent control
- Use rates vary
- Widespread resistance has been documented Worldwide
- Consult the label

Herbicide	Grasses Controlled
Clethodim	Annual grasses
Excel	Annual grasses including crabgrass
Venture	Annual grasses including crabgrass & Couch grass
Poast	Annual grasses including crabgrass & Couch grass

Prism (rimsulfuron)

- Group 2 – ALS inhibitor
- Use rate of 60 g ha⁻¹ - *use a surfactant*
- Can apply from emergence up to initiation of flowering
- Control of
 - Annual grasses
 - Mustards
 - Pigweed
- Suppression of
 - Lambsquarters
 - Corn spurrey



Preventative weed management

The potato vine crusher



Unprocessed control





Processed with PVC






Noxious species do not appear to persist in field margins & hedgerows



Weed ID & management resources



OMAFRA
Pest Manager



PMRA
Label Search



DropLeaf



Agriculture and
Agri-Food Canada

Agriculture et
Agroalimentaire Canada

Questions?

Charlottetown Weeds Lab

Peter Webb

Summer students

Nicolle MacDonald

Harrington Farm Crew

Andrew McKenzie-Gopsill, PhD

Research Scientist – Weed Science

Agriculture and Agri-Food Canada, Charlottetown

andrew.mckenzie-gopsill@agr.gc.ca (902) 314-3683